

## Calculus I

March 23-27

*Time Allotment: 40 minutes per day*

Student Name: \_\_\_\_\_

Teacher Name: \_\_\_\_\_

## Packet Overview

Date	Objective(s)	Page Number
Monday, March 23	Review of u-Substitution for Indefinite Integrals	2-6
Tuesday, March 24	u-Substitution change of variables review	7-10
Wednesday, March 25	Review of u-Substitution for Definite Integrals	11-13
Thursday, March 26	1. u-Substitution Practice 2. Quiz	14-16
Friday, March 27	1. Even and Odd Functions	17-19

**Additional Notes:** Each day's lessons will refer to textbook pages located at the end of this packet. Be sure to work all assigned example problems and exercises. For the exercises and quizzes, do your best to work the problem with a pencil, then check the solutions on the last page of your packet with a red pen.

Though not required to complete these assignments, Khan Academy's AP Calculus AB series of videos are a helpful resource for supplemental learning.

### Academic Honesty

I certify that I completed this assignment independently in accordance with the GHNO Academy Honor Code.

*Student signature:*

---

I certify that my student completed this assignment independently in accordance with the GHNO Academy Honor Code.

*Parent signature:*

---

## **Monday, March 23**

Calculus Unit: Integral Applications

Lesson 1: u-Substitution Review for Indefinite Integrals

### **Unit Overview: Integral Applications**

All of our hard work learning the theory and mechanics behind integrals is about to pay off! Our next unit will explore applications of integrals. Some of these applications include applying integrals to quantities in physics (position, velocity, acceleration, forces, work, energy, etc.) and calculating areas and volumes of geometric shapes, including deriving the volume formulas of cylinders, cones, and spheres. By the end of this unit, you will understand more deeply why the physics equations we use in physics class work. You will also never have to memorize another physics equation for the rest of physics I, nor will you have to remember area and volume equations because you will be able to use calculus to derive them!

**Objective:** Review u-substitution for definite integrals

1. Recall the steps required to evaluate integrals of composite functions.
2. Practice evaluating integrals using the method we learned earlier in the quarter.

### **Introduction to Lesson 1**

1. Turn to p. 295 in the textbook portion of your packet toward the back. Read carefully each paragraph and each step on p. 295-297.
2. As a warmup, see if you can evaluate the six integrals, a.-f. on the bottom of p. 295 (then check your answers on the key at the back of the packet).
3. Carefully work through Examples 1-3.
4. Do Exercises 1-6 on p. 304. Check answers with your red pen after you finish.
5. You may do work in the spaces that follow below, or you may do your work on a separate sheet of notebook paper. In either case, write the problem first, show all steps, and put your answer in a box.

p. 295

a)

b)

c)

d)

e)

f)

Example 1

Example 2

Example 3

p. 304

1)

2)

3)

4)

5)

6)

**Tuesday, March 24**

Calculus Unit: Applications of Integrals

Lesson 2: u-Substitution Review: Change of Variables

**Objective:** Be able to do this by the end of this lesson.

1. Evaluate more complicated integrals that require rewriting the integral in terms of  $u$  and  $du$ .

**Introduction to Lesson 2**

1. Read p. 298-300. Work Examples 4-7. Remember, you can always apply the Fundamental Theorem of Calculus to check your work by taking the derivative of your answer to see if you get back the original function back.
2. Do Exercises 7-14 on p. 304. Check answers in back of packet with a red pen!

Example 4

Example 5

Example 6

Example 7

p. 304  
7)



8)

9)

10)

11)

12)

13)

14)

**Wednesday, March 25**

Calculus Unit: Applications of Integrals

Lesson 3: u-Substitution Review: Definite Integrals

**Objective:** Be able to do this by the end of this lesson.

1. Evaluate definite integrals that require rewriting the integral in terms of  $u$  and  $du$  and changing the limits of integration.

**Introduction to Lesson 3**

1. Read p. 298-300. Work Examples 8-10.

2. Do Exercises 71-75 on p. 305. Check answers in back of packet with a red pen!

Example 8

Example 9

Example 10

p. 305  
71)

72)

73)

74)

75)

76)

**Thursday, March 26**

Calculus Unit: Applications of Integrals  
Lesson 4: u-Substitution Practice

**Objective:** Be able to do this by the end of this lesson.

1. Do a few warmup problems, check them, then take the quiz!

**Introduction to Lesson 4**

1. Do Exercises 15-18 on p. 304.

2. Do Exercises 43-45 and 63 and 64 on p. 305.

3. Take the quiz at the end of this section. Check answers with a red pen.

p. 304

15)

16)

17)

18)

p. 305  
63)

64)

Calculus I – Quiz on u-Substitution

Evaluate each integral.

1.  $\int (x + 2)^5 dx$

2.  $\int \sqrt{4t - 1} dt$

3.  $\int x^2 \sin(x^3) dx$

4.  $\int_{-1}^1 3x^2 \sqrt{x^3 + 1} dx$



**Friday, March 27**

Calculus Unit: Applications of Integrals

Lesson 4: Evaluating Integrals of Even and Odd Functions

**Objective:** Be able to do this by the end of this lesson.

1. Practicing evaluating definite integrals when you see the function is even or odd.

**Introduction to Lesson 4**

1. Reread p. 303. Write the steps for the Proof.
2. Do Exercises 101-104 and 106 (a)-(d) on p. 306.

p. 303 Proof

p. 306  
101)

102)

103)

104)

106 (a)

(b)

(c)

## Section 4.5

## Integration by Substitution

- Use pattern recognition to find an indefinite integral.
- Use a change of variables to find an indefinite integral.
- Use the General Power Rule for Integration to find an indefinite integral.
- Use a change of variables to evaluate a definite integral.
- Evaluate a definite integral involving an even or odd function.

## Pattern Recognition

In this section you will study techniques for integrating composite functions. The discussion is split into two parts—*pattern recognition* and *change of variables*. Both techniques involve a ***u*-substitution**. With pattern recognition you perform the substitution mentally, and with change of variables you write the substitution steps.

The role of substitution in integration is comparable to the role of the Chain Rule in differentiation. Recall that for differentiable functions given by  $y = F(u)$  and  $u = g(x)$ , the Chain Rule states that

$$\frac{d}{dx}[F(g(x))] = F'(g(x))g'(x).$$

From the definition of an antiderivative, it follows that

$$\begin{aligned}\int F'(g(x))g'(x) dx &= F(g(x)) + C \\ &= F(u) + C.\end{aligned}$$

These results are summarized in the following theorem.

**THEOREM 4.12** Antidifferentiation of a Composite Function

Let  $g$  be a function whose range is an interval  $I$ , and let  $f$  be a function that is continuous on  $I$ . If  $g$  is differentiable on its domain and  $F$  is an antiderivative of  $f$  on  $I$ , then

$$\int f(g(x))g'(x) dx = F(g(x)) + C.$$

If  $u = g(x)$ , then  $du = g'(x) dx$  and

$$\int f(u) du = F(u) + C.$$

**EXPLORATION**

**Recognizing Patterns** The integrand in each of the following integrals fits the pattern  $f(g(x))g'(x)$ . Identify the pattern and use the result to evaluate the integral.

$$\text{a. } \int 2x(x^2 + 1)^4 dx \quad \text{b. } \int 3x^2\sqrt{x^3 + 1} dx \quad \text{c. } \int \sec^2 x(\tan x + 3) dx$$


The next three integrals are similar to the first three. Show how you can multiply and divide by a constant to evaluate these integrals.

$$\text{d. } \int x(x^2 + 1)^4 dx \quad \text{e. } \int x^2\sqrt{x^3 + 1} dx \quad \text{f. } \int 2 \sec^2 x(\tan x + 3) dx$$

**NOTE** The statement of Theorem 4.12 doesn't tell how to distinguish between  $f(g(x))$  and  $g'(x)$  in the integrand. As you come more experienced at integration, your skill in doing this will increase. Of course, part of the key is familiarity with derivatives.

**STUDY TIP** There are several techniques for applying substitution, each differing slightly from the others. However, you should remember that the goal is the same with every technique—you are trying to find an antiderivative of the integrand.

Examples 1 and 2 show how to apply Theorem 4.12 *directly*, by recognizing the presence of  $f(g(x))$  and  $g'(x)$ . Note that the composite function in the integrand has an *outside function*  $f$  and an *inside function*  $g$ . Moreover, the derivative  $g'(x)$  is present as a factor of the integrand.



### EXAMPLE 1 Recognizing the $f(g(x))g'(x)$ Pattern

Find  $\int (x^2 + 1)^2(2x) dx$ .

**Solution** Letting  $g(x) = x^2 + 1$ , you obtain

$$g'(x) = 2x$$

and

$$f(g(x)) = f(x^2 + 1) = (x^2 + 1)^2.$$

From this, you can recognize that the integrand follows the  $f(g(x))g'(x)$  pattern. Using the Power Rule for Integration and Theorem 4.12, you can write

$$\int \overbrace{(x^2 + 1)^2(2x)}^{f(g(x))g'(x)} dx = \frac{1}{3}(x^2 + 1)^3 + C.$$

Try using the Chain Rule to check that the derivative of  $\frac{1}{3}(x^2 + 1)^3 + C$  is the integrand of the original integral.

### EXAMPLE 2 Recognizing the $f(g(x))g'(x)$ Pattern

Find  $\int 5 \cos 5x dx$ .

**Solution** Letting  $g(x) = 5x$ , you obtain

$$g'(x) = 5$$

and

$$f(g(x)) = f(5x) = \cos 5x.$$

From this, you can recognize that the integrand follows the  $f(g(x))g'(x)$  pattern. Using the Cosine Rule for Integration and Theorem 4.12, you can write

$$\int \overbrace{(\cos 5x)(5)}^{f(g(x))g'(x)} dx = \sin 5x + C.$$

You can check this by differentiating  $\sin 5x + C$  to obtain the original integrand.

**TECHNOLOGY** Try using a computer algebra system, such as *Maple*, *Derive*, *Mathematica*, *Mathcad*, or the *TI-89*, to solve the integrals given in Examples 1 and 2. Do you obtain the same antiderivatives that are listed in the examples?

The integrands in Examples 1 and 2 fit the  $f(g(x))g'(x)$  pattern exactly—you only had to recognize the pattern. You can extend this technique considerably with the Constant Multiple Rule

$$\int kf(x) dx = k \int f(x) dx.$$

Many integrands contain the essential part (the variable part) of  $g'(x)$  but are missing a constant multiple. In such cases, you can multiply and divide by the necessary constant multiple, as shown in Example 3.

### EXAMPLE 3 Multiplying and Dividing by a Constant

Find  $\int x(x^2 + 1)^2 dx$ .

**Solution** This is similar to the integral given in Example 1, except that the integrand is missing a factor of 2. Recognizing that  $2x$  is the derivative of  $x^2 + 1$ , you can let  $g(x) = x^2 + 1$  and supply the  $2x$  as follows.

$$\begin{aligned} \int x(x^2 + 1)^2 dx &= \int (x^2 + 1)^2 \left(\frac{1}{2}\right)(2x) dx && \text{Multiply and divide by 2.} \\ &= \frac{1}{2} \int \overbrace{(x^2 + 1)^2}^{f(g(x))} \overbrace{(2x)}^{g'(x)} dx && \text{Constant Multiple Rule} \\ &= \frac{1}{2} \left[ \frac{(x^2 + 1)^3}{3} \right] + C && \text{Integrate.} \\ &= \frac{1}{6} (x^2 + 1)^3 + C && \text{Simplify.} \end{aligned}$$

In practice, most people would not write as many steps as are shown in Example 3. For instance, you could evaluate the integral by simply writing

$$\begin{aligned} \int x(x^2 + 1)^2 dx &= \frac{1}{2} \int (x^2 + 1)^2 2x dx \\ &= \frac{1}{2} \left[ \frac{(x^2 + 1)^3}{3} \right] + C \\ &= \frac{1}{6} (x^2 + 1)^3 + C. \end{aligned}$$

**NOTE** Be sure you see that the *Constant Multiple Rule* applies only to *constants*. You cannot multiply and divide by a variable and then move the variable outside the integral sign. For instance,

$$\int (x^2 + 1)^2 dx \neq \frac{1}{2x} \int (x^2 + 1)^2 (2x) dx.$$

After all, if it were legitimate to move variable quantities outside the integral sign, you could move the entire integrand out and simplify the whole process. But the result would be incorrect.

## Change of Variables

With a formal **change of variables**, you completely rewrite the integral in terms of  $u$  and  $du$  (or any other convenient variable). Although this procedure can involve more written steps than the pattern recognition illustrated in Examples 1 to 3, it is useful for complicated integrands. The change of variable technique uses the Leibniz notation for the differential. That is, if  $u = g(x)$ , then  $du = g'(x) dx$ , and the integral in Theorem 4.12 takes the form

$$\int f(g(x))g'(x) dx = \int f(u) du = F(u) + C.$$

### EXAMPLE 4 Change of Variables

Find  $\int \sqrt{2x-1} dx$ .

**Solution** First, let  $u$  be the inner function,  $u = 2x - 1$ . Then calculate the differential  $du$  to be  $du = 2 dx$ . Now, using  $\sqrt{2x-1} = \sqrt{u}$  and  $dx = du/2$ , substitute to obtain

$$\begin{aligned} \int \sqrt{2x-1} dx &= \int \sqrt{u} \left( \frac{du}{2} \right) && \text{Integral in terms of } u \\ &= \frac{1}{2} \int u^{1/2} du && \text{Constant Multiple Rule} \\ &= \frac{1}{2} \left( \frac{u^{3/2}}{3/2} \right) + C && \text{Antiderivative in terms of } u \\ &= \frac{1}{3} u^{3/2} + C && \text{Simplify.} \\ &= \frac{1}{3} (2x-1)^{3/2} + C. && \text{Antiderivative in terms of } x \end{aligned}$$

**STUDY TIP** Because integration is usually more difficult than differentiation, you should always check your answer to an integration problem by differentiating. For instance, in Example 4 you should differentiate  $\frac{1}{3}(2x-1)^{3/2} + C$  to verify that you obtain the original integrand.



### EXAMPLE 5 Change of Variables

Find  $\int x\sqrt{2x-1} dx$ .

**Solution** As in the previous example, let  $u = 2x - 1$  and obtain  $dx = du/2$ . Because the integrand contains a factor of  $x$ , you must also solve for  $x$  in terms of  $u$ , as shown.

$$u = 2x - 1 \quad \Rightarrow \quad x = (u + 1)/2 \quad \text{Solve for } x \text{ in terms of } u.$$

Now, using substitution, you obtain

$$\begin{aligned} \int x\sqrt{2x-1} dx &= \int \left( \frac{u+1}{2} \right) u^{1/2} \left( \frac{du}{2} \right) \\ &= \frac{1}{4} \int (u^{3/2} + u^{1/2}) du \\ &= \frac{1}{4} \left( \frac{u^{5/2}}{5/2} + \frac{u^{3/2}}{3/2} \right) + C \\ &= \frac{1}{10} (2x-1)^{5/2} + \frac{1}{6} (2x-1)^{3/2} + C. \end{aligned}$$

To complete the change of variables in Example 5, you solved for  $x$  in terms of  $u$ . Sometimes this is very difficult. Fortunately it is not always necessary, as shown in the next example.

### EXAMPLE 6 Change of Variables

Find  $\int \sin^2 3x \cos 3x \, dx$ .

**Solution** Because  $\sin^2 3x = (\sin 3x)^2$ , you can let  $u = \sin 3x$ . Then

$$du = (\cos 3x)(3) \, dx.$$

Now, because  $\cos 3x \, dx$  is part of the original integral, you can write

$$\frac{du}{3} = \cos 3x \, dx.$$

Substituting  $u$  and  $du/3$  in the original integral yields

$$\begin{aligned} \int \sin^2 3x \cos 3x \, dx &= \int u^2 \frac{du}{3} \\ &= \frac{1}{3} \int u^2 \, du \\ &= \frac{1}{3} \left( \frac{u^3}{3} \right) + C \\ &= \frac{1}{9} \sin^3 3x + C. \end{aligned}$$

You can check this by differentiating.

$$\begin{aligned} \frac{d}{dx} \left[ \frac{1}{9} \sin^3 3x \right] &= \left( \frac{1}{9} \right) (3) (\sin 3x)^2 (\cos 3x) (3) \\ &= \sin^2 3x \cos 3x \end{aligned}$$

Because differentiation produces the original integrand, you know that you have obtained the correct antiderivative.

The steps used for integration by substitution are summarized in the following guidelines.

#### Guidelines for Making a Change of Variables

1. Choose a substitution  $u = g(x)$ . Usually, it is best to choose the *inner* part of a composite function, such as a quantity raised to a power.
2. Compute  $du = g'(x) \, dx$ .
3. Rewrite the integral in terms of the variable  $u$ .
4. Find the resulting integral in terms of  $u$ .
5. Replace  $u$  by  $g(x)$  to obtain an antiderivative in terms of  $x$ .
6. Check your answer by differentiating.

**STUDY TIP** When making a change of variables, be sure that your answer is written using the same variables as in the original integrand. For instance, in Example 6, you should not leave your answer as

$$\frac{1}{9} u^3 + C$$

but rather, replace  $u$  by  $\sin 3x$ .



## The General Power Rule for Integration

One of the most common  $u$ -substitutions involves quantities in the integrand that are raised to a power. Because of the importance of this type of substitution, it is given a special name—the **General Power Rule for Integration**. A proof of this rule follows directly from the (simple) Power Rule for Integration, together with Theorem 4.12.

### THEOREM 4.13 The General Power Rule for Integration

If  $g$  is a differentiable function of  $x$ , then

$$\int [g(x)]^n g'(x) dx = \frac{[g(x)]^{n+1}}{n+1} + C, \quad n \neq -1.$$

Equivalently, if  $u = g(x)$ , then

$$\int u^n du = \frac{u^{n+1}}{n+1} + C, \quad n \neq -1.$$

### EXAMPLE 7 Substitution and the General Power Rule

$$\text{a. } \int 3(3x-1)^4 dx = \int \overbrace{(3x-1)^4}^{u^4} \overbrace{(3)}^{du} dx = \frac{\overbrace{(3x-1)^5}^{u^5/5}}{5} + C$$

$$\text{b. } \int (2x+1)(x^2+x) dx = \int \overbrace{(x^2+x)^1}^{u^1} \overbrace{(2x+1)}^{du} dx = \frac{\overbrace{(x^2+x)^2}^{u^2/2}}{2} + C$$

$$\text{c. } \int 3x^2 \sqrt{x^3-2} dx = \int \overbrace{(x^3-2)^{1/2}}^{u^{1/2}} \overbrace{(3x^2)}^{du} dx = \frac{\overbrace{(x^3-2)^{3/2}}^{u^{3/2}/(3/2)}}{3/2} + C = \frac{2}{3}(x^3-2)^{3/2} + C$$

$$\text{d. } \int \frac{-4x}{(1-2x^2)^2} dx = \int \overbrace{(1-2x^2)^{-2}}^{u^{-2}} \overbrace{(-4x)}^{du} dx = \frac{\overbrace{(1-2x^2)^{-1}}^{u^{-1}/(-1)}}{-1} + C = -\frac{1}{1-2x^2} + C$$

$$\text{e. } \int \cos^2 x \sin x dx = - \int \overbrace{(\cos x)^2}^{u^2} \overbrace{(-\sin x)}^{du} dx = -\frac{\overbrace{(\cos x)^3}^{u^3/3}}{3} + C$$

### EXPLORATION

Suppose you were asked to find one of the following integrals. Which one would you choose? Explain your reasoning.

a.  $\int \sqrt{x^3+1} dx$  or

$$\int x^2 \sqrt{x^3+1} dx$$

b.  $\int \tan(3x) \sec^2(3x) dx$  or

$$\int \tan(3x) dx$$

Some integrals whose integrands involve quantities raised to powers cannot be found by the General Power Rule. Consider the two integrals

$$\int x(x^2+1)^2 dx \quad \text{and} \quad \int (x^2+1)^2 dx.$$

The substitution  $u = x^2 + 1$  works in the first integral but not in the second. In the second, the substitution fails because the integrand lacks the factor  $x$  needed for  $du$ . Fortunately, for this particular integral, you can expand the integrand as  $(x^2+1)^2 = x^4 + 2x^2 + 1$  and use the (simple) Power Rule to integrate each term.

### Change of Variables for Definite Integrals

When using  $u$ -substitution with a definite integral, it is often convenient to determine the limits of integration for the variable  $u$  rather than to convert the antiderivative back to the variable  $x$  and evaluate at the original limits. This change of variables is stated explicitly in the next theorem. The proof follows from Theorem 4.12 combined with the Fundamental Theorem of Calculus.

#### THEOREM 4.14 Change of Variables for Definite Integrals

If the function  $u = g(x)$  has a continuous derivative on the closed interval  $[a, b]$  and  $f$  is continuous on the range of  $g$ , then

$$\int_a^b f(g(x))g'(x) dx = \int_{g(a)}^{g(b)} f(u) du.$$

#### EXAMPLE 8 Change of Variables

Evaluate  $\int_0^1 x(x^2 + 1)^3 dx$ .

**Solution** To evaluate this integral, let  $u = x^2 + 1$ . Then, you obtain

$$u = x^2 + 1 \Rightarrow du = 2x dx.$$

Before substituting, determine the new upper and lower limits of integration.

Lower Limit	Upper Limit
When $x = 0$ , $u = 0^2 + 1 = 1$ .	When $x = 1$ , $u = 1^2 + 1 = 2$ .

Now, you can substitute to obtain

$$\begin{aligned} \int_0^1 x(x^2 + 1)^3 dx &= \frac{1}{2} \int_1^2 (x^2 + 1)^3 (2x) dx && \text{Integration limits for } x \\ &= \frac{1}{2} \int_1^2 u^3 du && \text{Integration limits for } u \\ &= \frac{1}{2} \left[ \frac{u^4}{4} \right]_1^2 \\ &= \frac{1}{2} \left( 4 - \frac{1}{4} \right) \\ &= \frac{15}{8}. \end{aligned}$$

Try rewriting the antiderivative  $\frac{1}{2}(u^4/4)$  in terms of the variable  $x$  and evaluate the definite integral at the original limits of integration, as shown.

$$\begin{aligned} \frac{1}{2} \left[ \frac{u^4}{4} \right]_1^2 &= \frac{1}{2} \left[ \frac{(x^2 + 1)^4}{4} \right]_0^1 \\ &= \frac{1}{2} \left( 4 - \frac{1}{4} \right) = \frac{15}{8} \end{aligned}$$

Notice that you obtain the same result.

**EXAMPLE 9** Change of Variables

Evaluate  $A = \int_1^5 \frac{x}{\sqrt{2x-1}} dx$ .

**Solution** To evaluate this integral, let  $u = \sqrt{2x-1}$ . Then, you obtain

$$u^2 = 2x - 1$$

$$u^2 + 1 = 2x$$

$$\frac{u^2 + 1}{2} = x$$

$$u \, du = dx.$$

Differentiate each side.


Before substituting, determine the new upper and lower limits of integration.

$$\begin{array}{l} \text{Lower Limit} \\ \text{When } x = 1, u = \sqrt{2-1} = 1. \end{array}$$

$$\begin{array}{l} \text{Upper Limit} \\ \text{When } x = 5, u = \sqrt{10-1} = 3. \end{array}$$


Now, substitute to obtain

$$\begin{aligned} \int_1^5 \frac{x}{\sqrt{2x-1}} dx &= \int_1^3 \frac{1}{u} \left( \frac{u^2 + 1}{2} \right) u \, du \\ &= \frac{1}{2} \int_1^3 (u^2 + 1) \, du \\ &= \frac{1}{2} \left[ \frac{u^3}{3} + u \right]_1^3 \\ &= \frac{1}{2} \left( 9 + 3 - \frac{1}{3} - 1 \right) \\ &= \frac{16}{3}. \end{aligned}$$



The region before substitution has an area of  $\frac{16}{3}$ .

**Figure 4.37**



The region after substitution has an area of  $\frac{16}{3}$ .

**Figure 4.38**

Geometrically, you can interpret the equation

$$\int_1^5 \frac{x}{\sqrt{2x-1}} dx = \int_1^3 \frac{u^2 + 1}{2} du$$

to mean that the two *different* regions shown in Figures 4.37 and 4.38 have the *same* area.

When evaluating definite integrals by substitution, it is possible for the upper limit of integration of the  $u$ -variable form to be smaller than the lower limit. If this happens, don't rearrange the limits. Simply evaluate as usual. For example, after substituting  $u = \sqrt{1-x}$  in the integral

$$\int_0^1 x^2(1-x)^{1/2} dx$$

you obtain  $u = \sqrt{1-x} = 0$  when  $x = 1$ , and  $u = \sqrt{1-0} = 1$  when  $x = 0$ . So the correct  $u$ -variable form of this integral is

$$-2 \int_1^0 (1-u^2)^2 u^2 du.$$

## Integration of Even and Odd Functions

Even with a change of variables, integration can be difficult. Occasionally, you can simplify the evaluation of a definite integral (over an interval that is symmetric about the y-axis or about the origin) by recognizing the integrand to be an even or odd function (see Figure 4.39).

### THEOREM 4.15 Integration of Even and Odd Functions

Let  $f$  be integrable on the closed interval  $[-a, a]$ .

1. If  $f$  is an *even* function, then  $\int_{-a}^a f(x) dx = 2 \int_0^a f(x) dx$ .
2. If  $f$  is an *odd* function, then  $\int_{-a}^a f(x) dx = 0$ .

**Proof** Because  $f$  is even, you know that  $f(x) = f(-x)$ . Using Theorem 4.12 with the substitution  $u = -x$  produces

$$\int_{-a}^0 f(x) dx = \int_a^0 f(-u)(-du) = -\int_a^0 f(u) du = \int_0^a f(u) du = \int_0^a f(x) dx.$$

Finally, using Theorem 4.6, you obtain

$$\begin{aligned} \int_{-a}^a f(x) dx &= \int_{-a}^0 f(x) dx + \int_0^a f(x) dx \\ &= \int_0^a f(x) dx + \int_0^a f(x) dx = 2 \int_0^a f(x) dx. \end{aligned}$$

This proves the first property. The proof of the second property is left to you (see Exercise 133).

### EXAMPLE 10 Integration of an Odd Function

Evaluate  $\int_{-\pi/2}^{\pi/2} (\sin^3 x \cos x + \sin x \cos x) dx$ .


**Solution** Letting  $f(x) = \sin^3 x \cos x + \sin x \cos x$  produces

$$\begin{aligned} f(-x) &= \sin^3(-x) \cos(-x) + \sin(-x) \cos(-x) \\ &= -\sin^3 x \cos x - \sin x \cos x = -f(x). \end{aligned}$$


So,  $f$  is an odd function, and because  $f$  is symmetric about the origin over  $[-\pi/2, \pi/2]$ , you can apply Theorem 4.15 to conclude that

$$\int_{-\pi/2}^{\pi/2} (\sin^3 x \cos x + \sin x \cos x) dx = 0.$$

**NOTE** From Figure 4.40 you can see that the two regions on either side of the y-axis have the same area. However, because one lies below the x-axis and one lies above it, integration produces a cancellation effect. (More will be said about this in Section 7.1.)




Even function



Odd function  
Figure 4.39

$$f(x) = \sin^3 x \cos x + \sin x \cos x$$



Because  $f$  is an odd function,

$$\int_{-\pi/2}^{\pi/2} f(x) dx = 0.$$

Figure 4.40

## Exercises for Section 4.5

In Exercises 1–6, complete the table by identifying  $u$  and  $du$  for the integral.

$$\int f(g(x))g'(x) dx \quad u = g(x) \quad du = g'(x) dx$$

1.  $\int (5x^2 + 1)^2(10x) dx$

2.  $\int x^2 \sqrt{x^3 + 1} dx$

3.  $\int \frac{x}{\sqrt{x^2 + 1}} dx$

4.  $\int \sec 2x \tan 2x dx$

5.  $\int \tan^2 x \sec^2 x dx$

6.  $\int \frac{\cos x}{\sin^2 x} dx$

In Exercises 7–34, find the indefinite integral and check the result by differentiation.

7.  $\int (1 + 2x)^4(2) dx$

8.  $\int (x^2 - 9)^3(2x) dx$

9.  $\int \sqrt{9 - x^2}(-2x) dx$

10.  $\int \sqrt[3]{1 - 2x^2}(-4x) dx$

11.  $\int x^3(x^4 + 3)^2 dx$

12.  $\int x^2(x^3 + 5)^4 dx$

13.  $\int x^2(x^3 - 1)^4 dx$

14.  $\int x(4x^2 + 3)^3 dx$

15.  $\int t\sqrt{t^2 + 2} dt$

16.  $\int t^3\sqrt{t^4 + 5} dt$

17.  $\int 5x\sqrt[3]{1 - x^2} dx$

18.  $\int u^2\sqrt{u^3 + 2} du$

19.  $\int \frac{x}{(1 - x^2)^3} dx$

20.  $\int \frac{x^3}{(1 + x^4)^2} dx$

21.  $\int \frac{x^2}{(1 + x^3)^2} dx$

22.  $\int \frac{x^2}{(16 - x^3)^2} dx$

23.  $\int \frac{x}{\sqrt{1 - x^2}} dx$

24.  $\int \frac{x^3}{\sqrt{1 + x^4}} dx$

25.  $\int \left(1 + \frac{1}{t}\right)^3 \left(\frac{1}{t^2}\right) dt$

26.  $\int \left[x^2 + \frac{1}{(3x)^2}\right] dx$

27.  $\int \frac{1}{\sqrt{2x}} dx$

28.  $\int \frac{1}{2\sqrt{x}} dx$

29.  $\int \frac{x^2 + 3x + 7}{\sqrt{x}} dx$

30.  $\int \frac{t + 2t^2}{\sqrt{t}} dt$

31.  $\int t^2 \left(t - \frac{2}{t}\right) dt$

32.  $\int \left(\frac{t^3}{3} + \frac{1}{4t^2}\right) dt$

33.  $\int (9 - y)\sqrt{y} dy$

34.  $\int 2\pi y(8 - y^{3/2}) dy$

In Exercises 35–38, solve the differential equation.

35.  $\frac{dy}{dx} = 4x + \frac{4x}{\sqrt{16 - x^2}}$


36.  $\frac{dy}{dx} = \frac{10x^2}{\sqrt{1 + x^3}}$

37.  $\frac{dy}{dx} = \frac{x + 1}{(x^2 + 2x - 3)^2}$


38.  $\frac{dy}{dx} = \frac{x - 4}{\sqrt{x^2 - 8x + 1}}$

**Slope Fields** In Exercises 39–42, a differential equation, a point, and a slope field are given. A *slope field* consists of line segments with slopes given by the differential equation. These line segments give a visual perspective of the directions of the solutions of the differential equation. (a) Sketch two approximate solutions of the differential equation on the slope field, one of which passes through the given point. (To print an enlarged copy of the graph, go to the website [www.mathgraphs.com](http://www.mathgraphs.com).) (b) Use integration to find the particular solution of the differential equation and use a graphing utility to graph the solution. Compare the result with the sketches in part (a).


39.  $\frac{dy}{dx} = x\sqrt{4 - x^2}$   
(2, 2)




40.  $\frac{dy}{dx} = x^2(x^3 - 1)^2$   
(1, 0)



41.  $\frac{dy}{dx} = x \cos x^2$   
(0, 1)



42.  $\frac{dy}{dx} = -2 \sec(2x) \tan(2x)$   
(0, -1)



In Exercises 43–56, find the indefinite integral.

43.  $\int \pi \sin \pi x \, dx$

44.  $\int 4x^3 \sin x^4 \, dx$

45.  $\int \sin 2x \, dx$

46.  $\int \cos 6x \, dx$

47.  $\int \frac{1}{\theta^2} \cos \frac{1}{\theta} \, d\theta$

48.  $\int x \sin x^2 \, dx$

49.  $\int \sin 2x \cos 2x \, dx$

50.  $\int \sec(1-x) \tan(1-x) \, dx$

51.  $\int \tan^4 x \sec^2 x \, dx$

52.  $\int \sqrt{\tan x} \sec^2 x \, dx$

53.  $\int \frac{\csc^2 x}{\cot^3 x} \, dx$

54.  $\int \frac{\sin x}{\cos^3 x} \, dx$

55.  $\int \cot^2 x \, dx$

56.  $\int \csc^2\left(\frac{x}{2}\right) \, dx$

In Exercises 57–62, find an equation for the function  $f$  that has the given derivative and whose graph passes through the given point.

Derivative	Point
57. $f'(x) = \cos \frac{x}{2}$	(0, 3)

58.  $f'(x) = \pi \sec \pi x \tan \pi x$

 $\left(\frac{1}{3}, 1\right)$ 

59.  $f'(x) = \sin 4x$

 $\left(\frac{\pi}{4}, -\frac{3}{4}\right)$ 

60.  $f'(x) = \sec^2(2x)$

 $\left(\frac{\pi}{2}, 2\right)$ 

61.  $f'(x) = 2x(4x^2 - 10)^2$

(2, 10)

62.  $f'(x) = -2x\sqrt{8-x^2}$

(2, 7)

In Exercises 63–70, find the indefinite integral by the method shown in Example 5.

63.  $\int x\sqrt{x+2} \, dx$

64.  $\int x\sqrt{2x+1} \, dx$

65.  $\int x^2\sqrt{1-x} \, dx$

66.  $\int (x+1)\sqrt{2-x} \, dx$

67.  $\int \frac{x^2-1}{\sqrt{2x-1}} \, dx$

68.  $\int \frac{2x+1}{\sqrt{x+4}} \, dx$

69.  $\int \frac{-x}{(x+1)\sqrt{x+1}} \, dx$

70.  $\int t\sqrt[3]{t-4} \, dt$

In Exercises 71–82, evaluate the definite integral. Use a graphing utility to verify your result.

71.  $\int_{-1}^1 x(x^2+1)^3 \, dx$

72.  $\int_{-2}^4 x^2(x^3+8)^2 \, dx$

73.  $\int_1^2 2x^2\sqrt{x^3+1} \, dx$

74.  $\int_0^1 x\sqrt{1-x^2} \, dx$

75.  $\int_0^4 \frac{1}{\sqrt{2x+1}} \, dx$

77.  $\int_1^9 \frac{1}{\sqrt{x}(1+\sqrt{x})^2} \, dx$

79.  $\int_1^2 (x-1)\sqrt{2-x} \, dx$

81.  $\int_0^{\pi/2} \cos\left(\frac{2x}{3}\right) \, dx$

82.  $\int_{\pi/3}^{\pi/2} (x+\cos x) \, dx$

76.  $\int_0^2 \frac{x}{\sqrt{1+2x^2}} \, dx$



78.  $\int_0^2 x\sqrt[3]{4+x^2} \, dx$

80.  $\int_1^5 \frac{x}{\sqrt{2x-1}} \, dx$

**Differential Equations** In Exercises 83–86, the graph of a function  $f$  is shown. Use the differential equation and the given point to find an equation of the function.



83.  $\frac{dy}{dx} = 18x^2(2x^3+1)^2$

84.  $\frac{dy}{dx} = \frac{-48}{(3x+5)^3}$



85.  $\frac{dy}{dx} = \frac{2x}{\sqrt{2x^2-1}}$



86.  $\frac{dy}{dx} = 4x + \frac{9x^2}{(3x^3+1)^{(3/2)}}$




In Exercises 87–92, find the area of the region. Use a graphing utility to verify your result.

87.  $\int_0^7 x\sqrt[3]{x+1} \, dx$


88.  $\int_{-2}^6 x^2\sqrt[3]{x+2} \, dx$




89.  $y = 2 \sin x + \sin 2x$




90.  $y = \sin x + \cos 2x$



91.  $\int_{\pi/2}^{2\pi/3} \sec^2\left(\frac{x}{2}\right) dx$



92.  $\int_{\pi/12}^{\pi/4} \csc 2x \cot 2x dx$



**✎** In Exercises 93–98, use a graphing utility to evaluate the integral. Graph the region whose area is given by the definite integral.

93.  $\int_0^4 \frac{x}{\sqrt{2x+1}} dx$

94.  $\int_0^2 x^3 \sqrt{x+2} dx$

95.  $\int_3^7 x \sqrt{x-3} dx$

96.  $\int_1^5 x^2 \sqrt{x-1} dx$

97.  $\int_0^3 \left( \theta + \cos \frac{\theta}{6} \right) d\theta$

98.  $\int_0^{\pi/2} \sin 2x dx$

**Writing** In Exercises 99 and 100, find the indefinite integral in two ways. Explain any difference in the forms of the answers.

99.  $\int (2x - 1)^2 dx$

100.  $\int \sin x \cos x dx$

In Exercises 101–104, evaluate the integral using the properties of even and odd functions as an aid.

101.  $\int_{-2}^2 x^2(x^2 + 1) dx$

102.  $\int_{-2}^2 x(x^2 + 1)^3 dx$

103.  $\int_{-\pi/2}^{\pi/2} \sin^2 x \cos x dx$

104.  $\int_{-\pi/2}^{\pi/2} \sin x \cos x dx$

105. Use  $\int_0^2 x^2 dx = \frac{8}{3}$  to evaluate each definite integral without using the Fundamental Theorem of Calculus.

(a)  $\int_{-2}^0 x^2 dx$

(b)  $\int_{-2}^2 x^2 dx$

(c)  $\int_0^2 -x^2 dx$

(d)  $\int_{-2}^0 3x^2 dx$

106. Use the symmetry of the graphs of the sine and cosine functions as an aid in evaluating each definite integral.

(a)  $\int_{-\pi/4}^{\pi/4} \sin x dx$

(b)  $\int_{-\pi/4}^{\pi/4} \cos x dx$

(c)  $\int_{-\pi/2}^{\pi/2} \cos x dx$

(d)  $\int_{-\pi/2}^{\pi/2} \sin x \cos x dx$

In Exercises 107 and 108, write the integral as the sum of the integral of an odd function and the integral of an even function. Use this simplification to evaluate the integral.

107.  $\int_{-4}^4 (x^3 + 6x^2 - 2x - 3) dx$

108.  $\int_{-\pi}^{\pi} (\sin 3x + \cos 3x) dx$

### Writing About Concepts

109. Describe why

$$\int x(5 - x^2)^3 dx \neq \int u^3 du$$

where  $u = 5 - x^2$ .

110. Without integrating, explain why

$$\int_{-2}^2 x(x^2 + 1)^2 dx = 0.$$

111. **Cash Flow** The rate of disbursement  $dQ/dt$  of a 2 million dollar federal grant is proportional to the square of  $100 - t$ . Time  $t$  is measured in days ( $0 \leq t \leq 100$ ), and  $Q$  is the amount that remains to be disbursed. Find the amount that remains to be disbursed after 50 days. Assume that all the money will be disbursed in 100 days.

112. **Depreciation** The rate of depreciation  $dV/dt$  of a machine is inversely proportional to the square of  $t + 1$ , where  $V$  is the value of the machine  $t$  years after it was purchased. The initial value of the machine was \$500,000, and its value decreased \$100,000 in the first year. Estimate its value after 4 years.

113. **Rainfall** The normal monthly rainfall at the Seattle-Tacoma airport can be approximated by the model

$$R = 3.121 + 2.399 \sin(0.524t + 1.377)$$

where  $R$  is measured in inches and  $t$  is the time in months, with  $t = 1$  corresponding to January. (Source: U.S. National Oceanic and Atmospheric Administration)

(a) Determine the extrema of the function over a one-year period.

(b) Use integration to approximate the normal annual rainfall. (Hint: Integrate over the interval  $[0, 12]$ .)

(c) Approximate the average monthly rainfall during the months of October, November, and December.



114. **Sales** The sales  $S$  (in thousands of units) of a seasonal product are given by the model

$$S = 74.50 + 43.75 \sin \frac{\pi t}{6}$$

where  $t$  is the time in months, with  $t = 1$  corresponding to January. Find the average sales for each time period.

- The first quarter ( $0 \leq t \leq 3$ )
- The second quarter ( $3 \leq t \leq 6$ )
- The entire year ( $0 \leq t \leq 12$ )

115. **Water Supply** A model for the flow rate of water at a pumping station on a given day is

$$R(t) = 53 + 7 \sin\left(\frac{\pi t}{6} + 3.6\right) + 9 \cos\left(\frac{\pi t}{12} + 8.9\right)$$

where  $0 \leq t \leq 24$ .  $R$  is the flow rate in thousands of gallons per hour, and  $t$  is the time in hours.

116. **Electricity** The oscillating current in an electrical circuit is

$$I = 2 \sin(60\pi t) + \cos(120\pi t)$$

where  $I$  is measured in amperes and  $t$  is measured in seconds. Find the average current for each time interval.

- $0 \leq t \leq \frac{1}{60}$
- $0 \leq t \leq \frac{1}{240}$
- $0 \leq t \leq \frac{1}{30}$

**Probability** In Exercises 117 and 118, the function

$$f(x) = kx^n(1-x)^m, \quad 0 \leq x \leq 1$$

where  $n > 0$ ,  $m > 0$ , and  $k$  is a constant, can be used to represent various probability distributions. If  $k$  is chosen such that

$$\int_0^1 f(x) dx = 1$$

the probability that  $x$  will fall between  $a$  and  $b$  ( $0 \leq a \leq b \leq 1$ ) is

$$P_{a,b} = \int_a^b f(x) dx.$$

117. The probability that a person will remember between  $a\%$  and  $b\%$  of material learned in an experiment is

$$P_{a,b} = \int_a^b \frac{15}{4} x \sqrt{1-x} dx$$

where  $x$  represents the percent remembered. (See figure.)

- For a randomly chosen individual, what is the probability that he or she will recall between 50% and 75% of the material?

- What is the median percent recall? That is, for what value of  $b$  is it true that the probability of recalling 0 to  $b$  is 0.5?





Figure for 117

118. The probability that ore samples taken from a region contain between  $a\%$  and  $b\%$  iron is

$$P_{a,b} = \int_a^b \frac{1155}{32} x^3(1-x)^{3/2} dx$$

where  $x$  represents the percent of iron. (See figure.) What is the probability that a sample will contain between

- 0% and 25% iron?
- 50% and 100% iron?




119. **Temperature** The temperature in degrees Fahrenheit in a house is

$$T = 72 + 12 \sin \left[ \frac{\pi(t-8)}{12} \right]$$

where  $t$  is time in hours, with  $t = 0$  representing midnight. The hourly cost of cooling a house is \$0.10 per degree.

- Find the cost  $C$  of cooling the house if its thermostat is set at 72°F by evaluating the integral

$$C = 0.1 \int_8^{20} \left[ 72 + 12 \sin \frac{\pi(t-8)}{12} - 72 \right] dt. \quad (\text{See figure.})$$






- (b) Find the savings from resetting the thermostat to 78°F by evaluating the integral

$$C = 0.1 \int_{10}^{18} \left[ 72 + 12 \sin \frac{\pi(t-8)}{12} - 78 \right] dt.$$

(See figure.)



120. **Manufacturing** A manufacturer of fertilizer finds that national sales of fertilizer follow the seasonal pattern

$$F = 100,000 \left[ 1 + \sin \frac{2\pi(t-60)}{365} \right]$$

where  $F$  is measured in pounds and  $t$  represents the time in days, with  $t = 1$  corresponding to January 1. The manufacturer wants to set up a schedule to produce a uniform amount of fertilizer each day. What should this amount be?

121. **Graphical Analysis** Consider the functions  $f$  and  $g$ , where

$$f(x) = 6 \sin x \cos^2 x \quad \text{and} \quad g(t) = \int_0^t f(x) dx.$$

- Use a graphing utility to graph  $f$  and  $g$  in the same viewing window.
- Explain why  $g$  is nonnegative.
- Identify the points on the graph of  $g$  that correspond to the extrema of  $f$ .
- Does each of the zeros of  $f$  correspond to an extremum of  $g$ ? Explain.
- Consider the function

$$h(t) = \int_{\pi/2}^t f(x) dx.$$

Use a graphing utility to graph  $h$ . What is the relationship between  $g$  and  $h$ ? Verify your conjecture.

122. Find  $\lim_{n \rightarrow +\infty} \sum_{i=1}^n \frac{\sin(i\pi/n)}{n}$  by evaluating an appropriate definite integral over the interval  $[0, 1]$ .
123. (a) Show that  $\int_0^1 x^2(1-x)^5 dx = \int_0^1 x^5(1-x)^2 dx$ .  
 (b) Show that  $\int_0^1 x^a(1-x)^b dx = \int_0^1 x^b(1-x)^a dx$ .
124. (a) Show that  $\int_0^{\pi/2} \sin^2 x dx = \int_0^{\pi/2} \cos^2 x dx$ .  
 (b) Show that  $\int_0^{\pi/2} \sin^n x dx = \int_0^{\pi/2} \cos^n x dx$ , where  $n$  is a positive integer.

**True or False?** In Exercises 125–130, determine whether the statement is true or false. If it is false, explain why or give an example that shows it is false.

125.  $\int (2x+1)^2 dx = \frac{1}{3}(2x+1)^3 + C$
126.  $\int x(x^2+1) dx = \frac{1}{2}x^2(\frac{1}{3}x^3+x) + C$
127.  $\int_{-10}^{10} (ax^3+bx^2+cx+d) dx = 2 \int_0^{10} (bx^2+d) dx$
128.  $\int_a^b \sin x dx = \int_a^{b+2\pi} \sin x dx$
129.  $4 \int \sin x \cos x dx = -\cos 2x + C$
130.  $\int \sin^2 2x \cos 2x dx = \frac{1}{3} \sin^3 2x + C$

131. Assume that  $f$  is continuous everywhere and that  $c$  is a constant. Show that

$$\int_{ca}^{cb} f(x) dx = c \int_a^b f(cx) dx.$$

132. (a) Verify that  $\sin u - u \cos u + C = \int u \sin u du$ .

(b) Use part (a) to show that  $\int_0^{\pi^2} \sin \sqrt{x} dx = 2\pi$ .

133. Complete the proof of Theorem 4.15.

134. Show that if  $f$  is continuous on the entire real number line, then

$$\int_a^b f(x+h) dx = \int_{a+h}^{b+h} f(x) dx.$$

### Putnam Exam Challenge

135. If  $a_0, a_1, \dots, a_n$  are real numbers satisfying

$$\frac{a_0}{1} + \frac{a_1}{2} + \dots + \frac{a_n}{n+1} = 0$$

show that the equation  $a_0 + a_1x + a_2x^2 + \dots + a_nx^n = 0$  has at least one real zero.

136. Find all the continuous positive functions  $f(x)$ , for  $0 \leq x \leq 1$ , such that

$$\int_0^1 f(x) dx = 1$$

$$\int_0^1 f(x)x dx = \alpha$$

$$\int_0^1 f(x)x^2 dx = \alpha^2$$

where  $\alpha$  is a real number.

These problems were composed by the Committee on the Putnam Prize Competition.  
 © The Mathematical Association of America. All rights reserved.

# Answer key for Calculus Packet 3-23 to 3-27

## Lesson 1

P. 295

a)  $\frac{1}{5}(x^2+1)^5 + C$    b)  $\frac{2}{3}(x^3+1)^{3/2} + C$    c)  $\frac{1}{2}\tan^2 x + 3\tan x + C$   
d)  $\frac{1}{10}(x^2+1)^5 + C$    e)  $\frac{2}{9}(x^3+1)^{3/2} + C$    f)  $\tan^2 x + 6\tan x + C$

P. 304

1)  $u = 5x^2 + 1$   
 $du = 10x dx$

2)  $u = x^3 + 1$   
 $du = 3x^2 dx$

3)  $u = x^2 + 1$   
 $du = 2x dx$

4)  $u = 2x$   
 $du = 2 dx$

5)  $u = \tan x$   
 $du = \sec^2 x dx$

6)  $u = \sin x$   
 $du = \cos x dx$

## Lesson 2

P. 304

7)  $\frac{(1+2x)^5}{5} + C$

8)  $\frac{(x^2-9)^4}{4} + C$

9)  $\frac{2}{3}(9-x^2)^{3/2} + C$

10)  $\frac{3}{4}(1-2x^2)^{4/3} + C$

11)  $\frac{(x^4+3)^3}{12} + C$

12)  $\frac{(x^3+5)^5}{15} + C$

13)  $\frac{1}{15}(x^3-1)^5 + C$

14)  $\frac{(4x^2+3)^4}{32} + C$

### Lesson 3

P. 305

71) 0

73)  $12 - \frac{8}{9}\sqrt{2}$

75) 2

72) 41,472

74)  $\frac{1}{3}$

### Lesson 4

P. 304

15)  $\frac{1}{3}(t^2+2)^{3/2} + C$

17)  $-\frac{15}{8}(1-x^2)^{4/3} + C$

16)  $\frac{1}{6}(t^4+5)^{3/2} + C$

18)  $\frac{2}{9}(u^3+2)^{3/2} + C$

43)  $-\cos(\pi x) + C$

45)  $-\frac{1}{2}\cos(2x) + C$

44)  $-\cos(x^4) + C$

\* Quiz key on separate page! \*

### Lesson 5

101)  $\frac{272}{15}$

103)  $\frac{2}{3}$

106) (a)  $\frac{8}{3}$

(c)  $-\frac{8}{3}$

(b)  $\frac{16}{3}$

(d) 8

102) 0

104) 0



## u-substitution Quiz Answers

$$1) \frac{(x+2)^6}{6} + C$$

$$2) \frac{1}{6} (4t-1)^{3/2} + C$$

$$3) -\frac{1}{3} \cos(x^3) + C$$

$$4) \frac{4\sqrt{2}}{3} \approx 1.89$$